

AUTISM & HYPERBARICS

The alarming number of children diagnosed with Autism Spectrum Disorders (ASD) in recent years has tragically increased; from 1 in 150 in 2000 to 1 in 88 by 2008. Due to that increase, more and more parents have turned to alternative forms of treatment to counteract the underlying symptoms of autism. Traditional medicines have often failed at producing desired results for many patients with autism and their families. However, studies have demonstrated hyperbaric oxygen therapy (HBOT) as an emerging treatment that parents and physicians are utilizing to help ameliorate the effects of ASD. This non-invasive treatment is as simple as a one hour daily session, where the parent and child enter a hyperbaric chamber and receive increased levels of oxygen. HBOT has been shown to decrease inflammation and improve cognitive functions, which have substantially helped thousands of ASD patients. Studies have demonstrated the benefits of HBOT for autism with the following:

Enhance Brain Healing & Repair with HBOT

- Increases Oxygen Delivery to Hypoxic Tissues
- Reduces Neuroinflammation
- Moderates Mitochondria Dysfunction
- Promotes Neurogenesis
- Stimulates Stem Cell Proliferation & Mobilization
- Reduces Seizure Frequency
- Actuates Neuroplasticity

Improve Physical Function with HBOT

- Promotes Receptive/Expressive Language
- Improves Movement & Walking
- Stimulates Better Eye Contact
- Improves Gross/Fine Motor Skills

Advance Mental Function with HBOT

- Improves Sensory/Cognitive Awareness
- Increases Attention Span & Alertness

Improve Behavioral Function with HBOT

- Improves Sleep Pattern
- Advances Social Skills & Interaction
- Reduces Aggression

Improve Digestive Function with HBOT

- Attenuates Gastrointestinal Issues
- Decreases Inflammation
- Enhances Nutritional Absorption



Scan QR Code to learn more and download related studies

Study: Autism Symptoms Ameliorated with HBOT

In 2009, a multi-center, randomized, controlled, double-blind trial was published, investigating the introduction of HBOT with 56 children diagnosed with ASD, ages 2-7. The study consisted of six separate HBOT centers that treated 30 children with 40 one hour sessions over the course of four weeks; whereas the remaining 26 children underwent sham treatment. Upon completion of this study, 80 percent of the treatment group's mean physician Clinical Global Impression (CGI) scores for overall functioning, receptive language, social interaction and eye contact significantly improved. Major improvements within the Aberrant Behavior Checklist (ABC) were observed in irritability, stereotypy, hyperactivity and speech. Finally, the treated children's ratings on the Autism Treatment Evaluation Checklist (ATEC) with respect to sensory/cognitive awareness, exemplified substantial improvement. Hyperbaric treatment was confirmed safe and well tolerated by the children and their parents in both groups.